



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

APRIL 6.

Mr. JOHN H. REDFIELD in the chair.

Twenty-five persons present.

Permission having been given, Prof. Heilprin withdrew his paper entitled "On a Giant *Conorbis* from the Oligocene of Florida."

The death of Charles Wilt, a member, was announced.

Fresh-water Sponges from Newfoundland: A new Species.—Mr. EDWARD POTTS stated that in the latter part of August, 1885, Mr. A. H. MacKay, of Pictou, Nova Scotia, whose success as a collector of fresh-water sponges in his own neighborhood has been already recorded (*Proc. Acad. Nat. Sci. Phila.*, 1884, p. 215, etc.), made a scientific visit to the island of Newfoundland. His explorations were mainly limited to the irregular peninsula of Avalon, the southeasterly extremity of the island, and the record of his collections beside mentioning the neighborhood of the city of St. Johns, embraces such familiar names as Trinity Bay, Harbor Grace, and Heart's Content, the landing place of the Atlantic cable.

He writes "I was extremely sorry that owing to my limited time and the impenetrability of the interior to any ordinary effort, I could not gain access to the great lakes in the heart and the western portion of the island. I have merely made a dip into a few of the ponds on the N. (?) E. coast." These are more particularly mentioned as Virginia and Ouidi Vidi Lakes, near St. Johns; Lady Lake, Bannerman Lake, Rocky Lake and Carbonear Lake, small bodies of water near Harbor Grace; and other lakelets and brooks upon the rocky ridges and near the sea level between Harbor Grace and Heart's Content. All this region is described as "the Canadian Huronian, the equivalent of the English Cambrian," and the collections were generally limited to the shallow margins of the ponds, where the sponges were found upon the under sides of splinters of hard slaty quartzites, in numbers very plentiful, but generally small; "from mere points to an inch or more in diameter."

It is to be regretted that the date of Mr. MacKay's visit was necessarily so early in the year, as the specimens collected were either immature or contained only the degenerate statoblasts of the preceding season. The information gathered as to the range

of temperature upon this island, is valuable as indicating so far the conditions of the growth of these and other organisms.

He says: "The Island is not extreme in its temperature, and the frost does not go very deep into the soil. The lakes freeze in November or December with ice at least a foot in thickness, and remain closed until the end of April. The average temperature during eight years, from 1857 to 1864, was 41.2° Fahr. Average maximum thermometer during the same time 83° , minimum 7° . In the year 1879, the mean temperature was 40.2° Fahr.; highest record August 3, 82° ; lowest December 22, $+4^{\circ}$. In Nova Scotia, though that is so much further south, the range is far greater, from $+96^{\circ}$ to -20° or -24° Fahr., with an annual average of 44° ."

The specimens of sponges so kindly forwarded by Mr. MacKay for examination and report, were more or less minute incrustations upon small stones, gathered as above indicated, and belonged to the species *Spongilla lacustris*, Auct.; *S. fragilis*, Leidy; *S. mackayi*, Carter; *Meyenia fluviatilis*, Auct., and *Heteromeyenia pictovens* and *Tubella pennsylvanica*, Potts. Of these, *Spongilla fragilis* was by far the most abundant, and our knowledge of its range is thus extended along the Eastern coast of North America from Florida to Newfoundland; whereas it had previously been traced westwardly to British Columbia near the Pacific Ocean, and more recently has been discovered in Russia, Bohemia and England. Beside the familiar species, *S. lacustris*, *S. fragilis* and *M. fluviatilis*, *Tubella pennsylvanica* has been rapidly enlarging its borders beyond the narrow limits of its original territorial designation; while *S. mackayi* and *H. pictovens* had previously been known only from the discoveries of Mr. MacKay in Nova Scotia.

One other form remains to be described, and at the suggestion of its discoverer it is hereby designated—


SPONGILLA NOVÆ TERRÆ, n. sp.

Sponge incrusting; sarcode of the young growth, a dense mass of minute spherical cells, embedding slender curving lines of fasciculated skeleton spicules, developing later into a very loose, open tissue, with few connecting spicules.

Gemmules rather numerous, unusually large, spherical; chitinous coat thin; "crust" apparently wanting.

Skeleton spicules relatively few, slender, cylindrical, smooth or sparsely microspined; gradually pointed.

Dermal or flesh spicules very abundant, minute *birotulates* of unequal size; shafts slender, cylindrical, occasionally spined; outer surface of rotules dome shaped; rays prolonged, terminations acute; malformations frequent. Mixed with occasional linear, spined spicules.



upon the crustless, chitinous body. Their shape when smooth is robust-fusiform, with pointed terminations; the great majority, however, have from one to six or more long spines, non-symmetrically placed, but with an evident tendency to group themselves at points about one-fourth the length of the spicule from one or both of its extremities.

Measurements: Diameter of gemmules 0.036 inches; skeleton spicules, 0.0068 by 0.0002 inches; length of average dermal spicule, 0.00066 inches; of gemmula spicule, 0.00145 inches.

Habitat.—Encrusting stones in shallow water.

Locality.—Lakes or ponds in the vicinity of Heart's Content, Newfoundland; collected by Mr. A. H. MacKay.

All the specimens of this sponge came from the neighborhood of Heart's Content, but whether they were gathered from a lake upon the heights or from a brook mentioned by Mr. MacKay near the sea level, does not seem entirely clear. The accompanying illustration (magnified 225 diameters) will suggest the peculiarities of its skeleton, dermal and gemmular spiculation. The striking resemblance of the dermal spicules to the minute birotulates, heretofore only known in a corresponding position, in the case of *Meyenia everetti*, will at once impress the student. These are however more variable in size, are occasionally spined, and have their rays more prolonged and more delicately terminated.

It is in the singular character of the spicules surrounding the gemmule that this sponge must attract peculiar attention. By the system of H. J. Carter, Esq., the fresh-water sponges are now classified into six genera, beside some conditional designations of forms in which the typical features are as yet undiscovered. These six, *Spongilla*, *Meyenia*, *Heteromeyenia*, *Tubella*, *Parmula* and *Carterius*, may again be associated into two groups, one of them including only the genus *Spongilla*, characterized by the *linear acerate* spicules surrounding the gemmule; and the other comprising all the other genera, where the spicule of corresponding significance is a *birotulate* or some easily recognized derivative of that type. Within this latter and larger group intermediate forms, connecting the defined genera, are frequent, and the location of species upon one side or other of

the distinctive line comparatively unimportant. Heretofore, between the genus *Spongilla* and those genera comprising the other group, there has been "a great gulf fixed." One only case in the past has suggested their possible association, or the development of one group from the other.

In *Meyenia acuminata*, Potts (Proc. Acad. Nat. Sci. Phila., 1882, p. 69), since regarded as a variety of *M. fluviatilis*, the shafts of the birotulates are prolonged at each extremity, forming acuminate terminations some distance beyond the surface of the rotules. In position, also, these spicules are abnormal, lying flat upon the chitinous coat, instead of resting upon one rotule, their shafts taking the position of radii, as is usual in this form. In fact we have the spicules of a *Meyenia* occupying the ordinary positions, and in degree approximating the forms of those peculiar to the *Spongillæ*.

In the present instance their intermediate character is still more striking, and while their form and position probably more closely associate them with the genus *Spongilla* in which the species has now been placed, the grouping of the ray-like species clearly suggests *Meyenia*. It has been an altogether unprecedented experience with the author to hesitate between these two genera, and it will be no cause of surprise if the future teacher shall shift it from its present position.

It must not escape notice that in both of these instances the gemmule are without "crust;"—that it is difficult to understand how birotulates could be supported in their ordinary positions *without* these embedding granules; and that we may not unreasonably infer that the change in position has induced the modification of type that we here find.

This collection of sponges, including the new species, has been examined coincidentally by H. J. Carter, Esq., F. R. S., and their identification and this description are believed to meet his approval.

APRIL 13.

Mr. JOHN H. REDFIELD in the chair.

Nineteen persons present.

The death of the Hon. John Welsh, a member, was announced.

APRIL 20.

Mr. GEO. W. TRYON, JR., in the chair.

Fifteen persons present.

The death of Andrew Nebinger, M.D., a member, was announced.